



University of Groningen

The Opacity of Spiral Galaxies from Counts of Distant Background Galaxies

Holwerda, B. W.; González, R. A.; Allen, R. J.; Kruit, P. C. van der

Published in:
Default journal

IMPORTANT NOTE: You are advised to consult the publisher's version (publisher's PDF) if you wish to cite from it. Please check the document version below.

Document Version
Publisher's PDF, also known as Version of record

Publication date:
2003

[Link to publication in University of Groningen/UMCG research database](#)

Citation for published version (APA):

Holwerda, B. W., González, R. A., Allen, R. J., & Kruit, P. C. V. D. (2003). The Opacity of Spiral Galaxies from Counts of Distant Background Galaxies. Default journal.

Copyright

Other than for strictly personal use, it is not permitted to download or to forward/distribute the text or part of it without the consent of the author(s) and/or copyright holder(s), unless the work is under an open content license (like Creative Commons).

Take-down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Downloaded from the University of Groningen/UMCG research database (Pure): <http://www.rug.nl/research/portal>. For technical reasons the number of authors shown on this cover page is limited to 10 maximum.

The Opacity of Spiral Galaxies from Counts of Distant Background Galaxies

B. W. Holwerda^{1,2}, R. A. González³, R. J. Allen¹ and
P. C. van der Kruit²

(1) *Space Telescope Science Institute, Baltimore, MD 21218, USA*

(2) *Kapteyn Institute, Landleven 12, 9747 AD Groningen, The Netherlands*

(3) *Centro de Radioastronomía y Astrofísica, Universidad Nacional Autónoma de México, 58190 Morelia, Michoacán, Mexico*

Abstract. We have applied the "Synthetic Field Method" on a sample of 20 nearby galaxies in order to determine the opacity of their disks. We present preliminary results on the radial dependence of cold dust absorption for 3 examples. The spirals NGC4535 and NGC4725 show significant absorption at a half-light radius. UGC2302, a LSB galaxy, shows much less opacity.

1. Introduction

The Synthetic Field Method (SFM) was developed by González et al. (1998) as a technique for determining the average opacity through the disk of a nearby spiral by counting more distant background galaxies. This number is compared with those of "synthetic fields", a known WFPc2 deep field, dimmed by a certain opacity, added to the foreground galaxy image.

González et al. (2003) examined the application of the method and concluded that the optimum results with HST/WFPc2 imaging would be obtained on Fornax and Virgo cluster galaxies, with the accuracy degrading for galaxies much closer or more distant.

Our sample was drawn from the HST archives, in part from the Cepheid Distance Scale HST Key project sample.

2. Results

We present preliminary results on NGC4535, NGC4725 and UGC2302. The numbers of real and simulated field galaxies were compared in annuli of deprojected radius.

2.1. NGC4535 and NGC4725

NGC4535 is a face-on SABc with a half-light radius of 5.1 kpc at a distance of 16 Mpc. Several spiral arms with bright HII regions are visible in the HST image. NGC4725 is a SABab ringed galaxy at 12 Mpc Its half-light radius is 5.4




Figure 1. The radial opacity measurement from field galaxy counts for NGC4535, NGC4725




Figure 2. The radial opacity measurement of UGC2302

kpc. The HST image shows much less structure. The radial opacity profiles of both however are similar (see Figure 1), showing 2.5 magnitudes of extinction in I at the half-light radius, dropping off to 0.5 magnitude.

2.2. UGC2302

UGC2302 is a LSB at 15 Mpc is a much more compact object. At comparable radii, it does not show as much absorption; less than a magnitude at 5 kpc from the center.

3. Conclusions

The uncertainties in these individual profiles are large due to low numbers of field galaxies and an added uncertainty due to their clustering. By averaging a number of fields we will improve on these uncertainties for an average radial opacity model of spiral galaxies. For different Hubble types, we intend to compare the radial dependence of opacity with the HI density profile to obtain cold dust-to-gas ratios.

References

- González, R. A., Allen, R. J., Dirsch, B., Ferguson, H. C., Calzetti, D., and Panagia, N. 1998, *ApJ* **506**, 152
- González, R. A., Loinard, L., Allen, R. J., and Muller, S. 2003, *AJ* **125**, 1182
- Holwerda, B. W., Allen, R. J., and van der Kruit, P. C.: 2002, *ASP Conf. Ser.* **273**: 337